

»MESSENGER RNA» IN EXPERIMENTAL GRANULOMAS

J. Ahonen and E. Kulonen

Department of Medical Chemistry, University of Turku, Finland

We have tried to identify messenger RNA specific for collagen by studying rapidly labelled RNA in experimental granulomas.

Nucleic acids were labelled *in vivo* and *in vitro* with ^{32}P -phosphate and ^3H -cytidine, and extracted by a modified phenolmethod, first at $+20^\circ\text{C}$. In order to release rapidly labelled RNA from nuclear-chromosomal material a second extraction was performed at $+65^\circ\text{C}$ (Georgiev and Mantieva, *Biochimiya* 27, 805, 1962). Ribosomal RNA together with rapidly labelled RNA was purified by salting-out with 3 *M* sodium acetate, and analyzed in sucrose-density-gradient ultracentrifugation and chromatography on methylated albumin-kieselguhr (MAK). The distribution of radioactive phosphate in the nucleotides was determined after an alkaline hydrolysis. Collagen-producing granulomas (18 day old) were compared with those which were not yet producing collagen (6 day old).

The following features were found in collagen-producing granulomas: (1) a RNA fraction of 22S which was labelled most rapidly; (2) the rapidly-labelled RNA was extracted efficiently at $+65^\circ\text{C}$ and it was eluted in MAK-chromatography at a relatively low salt concentration which indicates high G- and C-content. The distribution of ^{32}P among the nucleotides of this fraction was very uneven.